

## 2013-2015 LA HOPE Surveys

Domain/Subdomain		<i>Nutrition &amp; Physical Activity / Obesity and Overweight</i>			<i>Nutrition &amp; Physical Activity / Folic Acid Supplementation</i>			<i>Nutrition &amp; Physical Activity / Exercise/Physical Activity</i>		
Indicator Description		Percentage of women having an infant/fetal death who are overweight* or obese** based on body mass index (BMI) *Overweight: percentage of women with a pre-pregnancy BMI $\geq$ 25 kg/m2 but < 30 kg/m2 **Obesity: percentage of women with a pre-pregnancy BMI $\geq$ 30 kg/m2			Percentage of women having an infant/fetal death who did not take a multivitamin, prenatal vitamin, or a folic acid supplement during one month prior to pregnancy.			Percentage of women having an infant/fetal death who did not exercise during pregnancy.		
		%	95% CI		%	95% CI		%	95% CI	
<b>LA County</b>		59.8	56.1	63.4	46.0	42.4	49.5	43.7	39.9	47.4
<b>Race/Ethnicity</b>	<b>White</b>	43.2	34.1	52.2	28.7	20.6	36.8	33.5	24.5	42.5
	<b>Latina</b>	66.6	62.1	71.1	52.8	48.3	57.3	42.9	38.3	47.5
	<b>African-American</b>	67.6	57.8	77.5	42.7	32.9	52.6	48.5	38.2	58.8
	<b>Asian/Pacific Islander</b>	30.0	15.5	44.6	28.9	14.8	43.0	54.1	37.2	71.0
<b>Service Planning Areas</b>	<b>SPA 1 Antelope Valley</b>	73.9	62.8	84.9	64.0	51.0	77.0	59.1	45.1	73.1
	<b>SPA 2 San Fernando</b>	57.8	50.3	65.3	44.7	37.3	52.0	44.2	36.5	51.9
	<b>SPA 3 San Gabriel</b>	65.5	55.9	75.1	41.5	32.0	51.1	41.6	31.1	52.0
	<b>SPA 4 Metro</b>	41.8	29.7	53.9	40.3	28.5	52.1	42.8	30.0	55.6
	<b>SPA 5 West</b>	22.1*	8.3	35.9	24.4*	10.6	38.1	40.0	21.8	58.2
	<b>SPA 6 South</b>	70.2	62.2	78.2	50.6	42.6	58.7	42.7	34.3	51.1
	<b>SPA 7 East</b>	69.0	60.1	77.8	52.6	43.2	62.0	40.7	31.1	50.3
	<b>SPA 8 South Bay</b>	49.4	38.6	60.2	41.4	31.0	51.8	44.2	33.3	55.1

\*The estimate is statistically unstable (relative standard error > 25%) and therefore may not be appropriate to use for planning or policy purposes.

[For more information about LA HOPE project please click here](#)