

## 2007-2009 LA HOPE surveys

Domain/Subdomain		<i>Nutrition &amp; Physical Activity / Obesity and Overweight</i>			<i>Nutrition &amp; Physical Activity / Folic Acid Supplementation</i>			<i>Nutrition &amp; Physical Activity / Exercise/Physical Activity</i>			
Indicator Description		Percentage of women having an infant/fetal death who are overweight* or obese** based on body mass index (BMI) *Overweight: percentage of women with a pre-pregnancy BMI $\geq$ 25 kg/m <sup>2</sup> but < 30 kg/m <sup>2</sup> **Obesity: percentage of women with a pre-pregnancy BMI $\geq$ 30 kg/m <sup>2</sup>			Percentage of women having an infant/fetal death who did not take a multivitamin, prenatal vitamin, or a folic acid supplement during one month prior to pregnancy.			Percentage of women having an infant/fetal death who did not exercise during pregnancy.			
		%	95% CI		%	95% CI		%	95% CI		
<b>LA County</b>		54.6	51.1	58.1	60.2	56.8	63.6	46.7	43.2	50.3	
<b>Race/Ethnicity</b>	<b>White</b>	35.2	27.1	43.2	38.5	30.4	46.6	40.7	32.1	49.4	
	<b>Latina</b>	59.4	54.9	64.0	67.0	62.8	71.1	49.2	44.6	53.7	
	<b>African-American</b>	62.5	54.5	70.6	66.0	58.5	73.6	45.8	37.6	53.9	
	<b>Asian/Pacific Islander</b>	42.1	25.9	58.2	43.1	27.8	58.4	36.6	20.5	52.7	
<b>Service Planning Areas</b>	SPA 1	Antelope Valley	62.8	51.3	74.2	63.0	52.0	74.0	56.8	45.3	68.3
	SPA 2	San Fernando	51.2	42.9	59.5	49.8	41.6	58.0	47.8	39.3	56.2
	SPA 3	San Gabriel	49.7	40.7	58.7	56.8	48.3	65.4	49.6	40.3	58.9
	SPA 4	Metro	49.7	38.7	60.6	62.0	51.9	72.1	36.2	26.1	46.2
	SPA 5	West	17.3*	5.4	29.2	32.6	17.9	47.3	32.5*	16.2	48.9
	SPA 6	South	64.4	56.7	72.2	73.6	66.8	80.3	45.6	37.7	53.4
	SPA 7	East	61.9	52.3	71.5	58.4	48.6	68.2	50.4	40.6	60.1
	SPA 8	South Bay	53.4	43.8	63.0	65.7	56.9	74.5	46.5	36.8	56.3

\*The estimate is statistically unstable (relative standard error > 25%) and therefore may not be appropriate to use for planning or policy purposes.

[For more information about LA HOPE project please click here](#)